

## The combined effects of a monotonous diet and exposure to thiamethoxam on the performance of bumblebee micro-colonies

Article (Accepted Version)

Dance, C, Botías, C and Goulson, D (2017) The combined effects of a monotonous diet and exposure to thiamethoxam on the performance of bumblebee micro-colonies. *Ecotoxicology and Environmental Safety*, 139. pp. 194-201. ISSN 0147-6513

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/66676/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

The combined effects of a monotonous diet and exposure to thiamethoxam on the performance of bumblebee micro-colonies

Dance, C., Botías, C., & Goulson, D.

School of Life Sciences, University of Sussex, Falmer, Brighton, BN1 9QG

Running title: Combined effects of diet and pesticides on bumblebees.

Key words: Bees; neonicotinoids; pollination; pollen quality; stressors; bee health

## Abstract

There is a pressing need to better understand the factors contributing to declines of wild pollinators such as bumblebees. Many different contributors have been postulated including: loss of flower-rich habitats and nesting sites; monotonous diets; impacts of invasive pathogens; exposure to pesticides such as neonicotinoids. Past research has tended to investigate the impacts of these stressors in isolation, despite the increasing recognition that bees are simultaneously exposed to a combination of stressors, with potentially additive or synergistic effects. No studies to date have investigated the combined effects of a monotonous diet and exposure to pesticides. Using queenless micro-colonies of *Bombus terrestris audax*, we examined this interaction by providing bees with monofloral or polyfloral pollen that was either contaminated with field-realistic levels of thiamethoxam, a commonly used neonicotinoid, or not contaminated. Both treatments were found to have a significant effect on various parameters relating to micro-colony performance. Specifically, both pesticide-treated micro-colonies and those fed monofloral pollen grew more slowly than those given polyfloral pollen or pollen without pesticides. The two factors appeared to act additively. Micro-colonies given monofloral pollens also exhibited lower reproductive efforts and produced smaller drones. Although further research is needed to examine whether similar effects are found in whole colonies, these findings increase our understanding of the likely effects of multiple stressors associated with agricultural intensification on bee declines.

## Introduction

Considering the invaluable ecosystem services provided by bees, particularly through their pollination of wildflowers and crops (Gallai et al., 2009), emerging evidence for declines of some species are a great cause for concern. For wild bees, evidence of decline is most clear in bumblebees (Rasmont et al., 2005; Biesmeijer et al., 2006; Kosior et al., 2007; Goulson et al., 2008; Xie et al., 2008; Grixti et al., 2009; Williams & Osborne, 2009; Cameron et al., 2011; Goulson et al. 2015)

Many factors have been implicated in contributing to worldwide losses in pollinator stocks, the most prominent of which are habitat loss and degradation, exposure to harmful agrochemicals such as pesticides, competition from invasive species, pathogens and parasites and diet stress, and climate change is only likely to further exacerbate these existing pressures (Brown & Paxton, 2009; Potts et al., 2010; Goulson et al., 2015). Generally recognised as the most significant driver of declines in biodiversity at a global scale is land-use change and its concomitant habitat loss (Foley et al., 2005), and the same is true for losses of bees (Goulson et al., 2008; Brown & Paxton, 2009; Potts et al., 2010; Winfree, 2010; Goulson et al., 2015). As increasing amounts of natural, flower-rich habitat is converted to agricultural land, the availability of suitable, undisturbed nesting sites and consistent and varied floral resources, on which many species of wild bee depend, is reduced (Carvell, 2002; Williams & Osborne, 2009; Goulson et al. 2015). For example, the range and abundance of many plants on which bumblebees tend to forage have declined in the United Kingdom (Carvell et al., 2006; Kleijn & Raemakers, 2008), with 97% of flower-rich grasslands having been lost in Britain in the 20<sup>th</sup> century (Howard et al., 2003; Goulson et al., 2015). Often what is left is a more homogenous landscape, characterised by short, temporally and spatially isolated blooming periods of mass-flowering crops such as oilseed rape and canola (Westpal et al., 2006; Osborne et al., 2008). These landscapes are generally less suited to pollinators; in a meta-analysis of 54 studies, Winfree et al. (2009) found habitat loss to be the most significant contributor to losses in wild bee richness and abundance. Similarly, Ricketts et al. (2008), in a review of 23 studies detected a negative correlation between wild bee diversity and distance from areas of natural habitat.

Due to these losses in the extent of wildflowers, it has been proposed that mass-flowering crops could provide valuable resources for pollinators (Westphal et al., 2003). However, as they are only available for such short period of time, they might not be sufficient to sustain viable pollinator populations (Kremen et al., 2007). Furthermore, bees inhabiting areas of intensive farmland will almost certainly have more monotonous diets than they would have done in their evolutionary past (Goulson et al., 2015) and this has caused concern that pollinators may be adversely affected by inadequate nutrition, although the effects of diet stress have been little investigated. It is well known that the nutritive quality of both pollen and nectar of different plants is highly variable (Hanley et al., 2008). For example, pollen protein content can range from 2.5 to 61% (Roulston et al.,

2000). Therefore, it is not surprising that pollen diet can have important implications for the development of bee colonies. One study examining the effects of pollen quality and diversity on honey bees found that bee physiology and immune system function were both increased when pollen diet was of higher quality (i.e. higher protein content) and more diverse (i.e. polyfloral; pollen originating from multiple plant species) (Di Pasquale et al., 2013). Furthermore, studies on bumblebees have also indicated the importance of pollen diet in colony development and brood production, the general trends being that colonies perform better when pollen source is varied or of higher quality (Génissel et al., 2002; Tasei & Aupinel, 2008a; Vanderplanck et al., 2014; Baloglu & Gurel, 2015; Moerman et al., 2015). Whilst these studies have primarily been intended for maximising the efficiency of commercial bumblebee rearing for crop pollination, they nevertheless may help in the understanding of the influence of agricultural intensification on bee health and nutrition (Di Pasquale et al., 2013).

Not only does agricultural intensification lower the availability of suitable habitats and food sources, remaining habitats may be further degraded due to the use of agrochemicals, such as herbicides, fungicides and insecticides, many of which are toxic to pollinators (Williams & Osborne, 2009; Goulson et al., 2015). Of the pesticides to which bees are likely to be exposed, neonicotinoids have attracted most attention and debate. Since their development in the 1980s and their commercial availability in the 1990s (Kollmeyer et al., 1999), they have rapidly become the most widely used class of insecticides in the world (Goulson, 2013). As nicotinic acetylcholine receptor (nAChR) agonists, they bind to receptors in the central nervous system (Elbert et al., 2008). In low concentrations, this causes nervous stimulation but higher doses can lead to paralysis and death. Their water solubility and systemic nature means that they are readily absorbed by roots and leaves and transported around the whole plant protecting all the plant tissues. This however has important implications for pollinators as varying concentrations of these chemicals are often found in the pollen and nectar of both treated crops and nearby wildflowers (Botías et al., 2015). Whilst the concentrations of neonicotinoids are generally not sufficient to cause rapid mortality in pollinators (Goulson, 2013), a wide range of sub-lethal effects have been documented including reductions in foraging and homing abilities, (Yang et al., 2008; Schneider et al., 2012), weakened immune function (Di Prisco et al., 2013), reduced food consumption (Tasei et al., 2000) reduced nest growth, and lower reproductive capacity (Gill et al., 2012; Laycock et al., 2012; Whitehorn et al., 2012). The majority of the controversy over the effects of neonicotinoids has been concerned with whether bees actually encounter large enough amounts in the wild to cause them significant harm (Godfray et al., 2014), and this may in part be down to the huge variability in concentrations of these chemicals found in the field (Blacquière et al., 2012). However, recent studies have shown that

101 persistence of neonicotinoids in untreated wildflowers means that exposure is likely to be more  
102 extensive than previously thought (Botías et al., 2015).

103 The majority of studies to date have focussed on the impacts of imidacloprid on bees, but  
104 other neonicotinoids such as thiamethoxam and clothianidin are now used more frequently (Laycock  
105 et al., 2014). Whilst detrimental effects of thiamethoxam to honey bees and bumblebees have been  
106 documented at fairly high doses, ranging from 67 ng/g to higher than 100 ng/g (Mommaerts et al.,  
107 2010; Henry et al., 2012), residues in crops and wildflowers do not tend to reach these levels and are  
108 more often in the range of 1 to 12 ng/g (Arnold et al., 2012; Dively & Kamel, 2012; Stoner & Eitzer,  
109 2012; Botías et al., 2015). Evidence of effects at field-realistic levels on bumblebees is conflicting;  
110 Elston et al. (2013) detected a significant reduction in nest building and brood production at levels as  
111 low as 1 ng/g and 10 ng/g respectively, whilst others found no effects with doses of 10 ng/g  
112 (Mommaerts et al., 2010; Laycock et al., 2014). This discrepancy may be in part explained by  
113 differences in the methodologies of the studies. The two latter studies only exposed bees to  
114 thiamethoxam in dietary syrup and not pollen (Mommaerts et al., 2010; Laycock et al., 2014),  
115 despite the fact that neonicotinoids are present in both pollen and nectar. Most recently, Goulson  
116 (2014) found that concentrations of thiamethoxam in pollen stores of free-flying bumblebee nests in  
117 the range 0 to 1.6 ppb strongly and negatively correlated with colony performance, but these nests  
118 were also exposed to a cocktail of other neonicotinoids so disentangling effects of particular  
119 compounds is difficult.

120 The majority of scientific literature and public debate on the topic of bee health has tended  
121 to focus on the impacts of the individual drivers of pollinator declines in isolation, with the emphasis  
122 often on attempting to identify the sole or primary cause of bee declines (Potts et al., 2010; Goulson  
123 et al., 2015). However, it has been increasingly recognised that these drivers rarely act in isolation,  
124 and that in the wild, bees will commonly be faced by combinations of numerous different stressors  
125 that may interact additively or synergistically (Potts et al., 2010; Goulson et al., 2015). These kinds of  
126 interactions have been documented between different agrochemicals, whereby chemicals such as EBI  
127 fungicides greatly increase the toxicity of insecticides (Pilling & Jepson, 1993; Schmuck et al., 2003;  
128 Sgolastra et al., 2016). Furthermore, there is increasing evidence that exposure to pesticides can  
129 lower immune system function, making bees more susceptible to damage from pathogens, such as  
130 *Nosema ceranae* (Alaux et al., 2010; James & Xu, 2012; Pettis et al., 2012; Di Prisco et al., 2013). Diet  
131 stress has also been implicated in affecting the ability of bumblebees to fight off infection from a  
132 trypanosome parasite, with starved bees experiencing much higher mortality rates (Brown et al.,  
133 2000). Moreover, a recent study found that the combined exposure to poor quality pollen and the  
134 neonicotinoid thiamethoxam had detrimental effects on hypopharyngeal gland development of  
135 honeybees (Renzi et al., 2016). It has thus been hypothesised that nutritional stress may have the

potential to lower bees' capacity to withstand the effects of pesticides (Goulson et al., 2015) although this has yet to be tested in bumblebees.

Here we investigate the combined effects of a monotonous diet and exposure to thiamethoxam on bumblebee micro-colonies. Queenless micro-colonies are considered to be reliable indicators of trends in larger queenright colonies (Tasei & Aupinel, 2008b), and are recommended for risk assessments of agrochemicals by the European Food Safety Authority (EFSA, 2013). Monotonous and varied diets were simulated by feeding micro-colonies either monofloral or polyfloral diets. Simultaneously, half the micro-colonies in each diet treatment were also exposed to environmentally-realistic levels of thiamethoxam in both pollen and syrup over a period of 17 days, after which point uncontaminated pollen and syrup were provided. Colonies were observed and performance parameters were recorded both during and after exposure to determine the effects of the two treatments and their interaction on measures of colony performance.

## Methods

Honeybee-collected *Cistus* spp. pollen was purchased from Pollenergie® (France) and a honeybee-collected polyfloral pollen blend was purchased from Biobest (Belgium) via Agralan Ltd (Swindon, UK). As honeybee pollen loads can potentially contain viable *Nosema ceranae* spores (Higes et al., 2008), deformed wing virus (Singh et al., 2010) and other bee pathogens (Graystock et al., 2016), all the pollen provided to our micro-colonies was sterilized to exclude honeybee pathogen spill-over effects. Polyfloral pollen was sterilised by Biobest using gamma irradiation with a cobalt-60 source at dose rates between 25-45 kGy. We were unable to use this approach so monofloral pollen was sterilised by a 30 minutes cycle exposure to ultraviolet germicidal light (254 nm). Whilst a single study has indicated that gamma irradiation has no effect on pollen protein content (Junjie et al., 1998), it is possible that the different sterilisation methods may have affected some other nutritive quality of pollen. However, the effect of sterilisation techniques on pollen quality has been largely unexplored.

Protein content of both the monofloral and polyfloral pollens was calculated to be 10.15 and 12.60%, respectively, using the Bradford method (Bradford, 1976), adapted for use with the NanoDrop 2000/2000c (Thermo Scientific, 2010). The polyfloral pollen was examined using a microscope and four dominant types of approximately equal representation were identified (Asteraceae *Taraxacum* type, 23.4%; Rosaceae *Rubus* type, 20.3%; Rosaceae *Crataegus/Malus* type, 18.6%; Papaveraceae *Papaver* type, 14.9%). The remaining 22.8% was made up of 7 more pollen types, each representing less than 5% of the total volume. *Cistus* pollen was added to the polyfloral blend so that it was at a similar proportion to the 5 main pollen groups. This was in order to

minimise any detrimental or favourable effects of any toxin and/or additional nutrient present in *Cistus* pollen acting only on monofloral treatment colonies.

Four colonies of *Bombus terrestris audax*, each with approximately 100 workers, were purchased from Biobest (Belgium) via Agralan Ltd (Swindon, UK). Forty queenless micro-colonies were established by placing 5 workers from one of the four queenright colonies into circular plastic boxes (diameter 11cm, height 9 cm) with an aluminium mesh cover to allow air ventilation. Micro-colonies were kept in a dark room with controlled conditions throughout the entire study period ( $50 \pm 5\%$  humidity and  $24 \pm 1^\circ\text{C}$ ). Workers were left for 2 days to acclimatise to their new environment, during which time uncontaminated polyfloral pollen and syrup were supplied *ad libitum*. After the 2 days, micro-colonies were assigned to one of four treatment groups with 10 micro-colonies per group. Micro-colonies with workers originating from each queenright colony were assigned evenly to the four different treatment groups in order to control for effects of the workers' colony of origin on performance. Micro-colonies were weighed and a small amount of wax from the corresponding queenright colony was then added to stimulate oviposition. Half of the micro-colonies received monofloral pollen and half were given polyfloral pollen. All groups received the same inverted sugar syrup solution (50% Ambrosia syrup, EH Thorne Ltd), and groups were supplied with their particular pollen diet throughout the 5 week study period. Within each diet treatment, half of the colonies were exposed to thiamethoxam and the others were provided with uncontaminated food. Pollen and nectar were dosed with thiamethoxam at field realistic levels of 3.5 ppb (Botías et al. 2015). The period of exposure to thiamethoxam lasted 17 days, after which time, all groups were supplied with uncontaminated pollen and nectar.

Colonies were observed and performance parameters including worker mortality, micro-colony growth, reproductive effort and food collection were recorded. Daily observations consisted of counting and removing any dead workers or newly emerged males. Males were weighed and their thoraxes were measured using callipers. Their lipid content was measured using a protocol slightly modified from Brown et al. (2000). Briefly, the whole body of each bumblebee was dried at  $70^\circ\text{C}$  for 5 days and weighed on a precision balance. Every dried bee was then placed in an Eppendorf tube containing 1 ml diethyl ether for 24 h to dissolve lipids, vortexing the tubes for 30 seconds every 3-4 hours (except for the overnight period). The bees were then rinsed in fresh diethyl ether, and subsequently dried at  $70^\circ\text{C}$  for a further 5 days and finally reweighed. The amount of fat in each bumblebee was taken from the difference between the first and second weight measurements.

Every three days, syrup and pollen feeders were weighed to measure collection and fresh pollen and syrup were provided. Data on food collection were also used to calculate the average amount of active compound collected by each bee. We consider this pollen and syrup collection rather than consumption as some syrup was stored in nectar pots and pollen was used to provision

brood. Five identical plastic boxes to those used for the bee micro-colonies were kept with full syrup feeders and weighed every 3 days in order to control for any effects of evaporation in syrup collection analyses. The micro-colonies were also weighed and the number of brood cells and nectar pots was noted.

At the end of the fifth week, all the colonies were frozen and dissected. The numbers of larvae and pupae were counted and the workers were weighed.

All statistical analyses were carried out using SPSS 21.0. Data were first tested for normality using a Shapiro-Wilk test. Where data were normally distributed, generalized linear models (GLM) were used to test for effects of pollen diet and exposure to pesticides, and any interactions between the two, on the colony performance parameters. Where distributions were not normal (e.g. numbers of males produced), non-parametric tests were used. Analyses were also carried out to determine whether there were any significant differences between colony growth and the food collection of pesticide treated groups during the period of exposure and after the period of exposure. Two linear regressions were calculated for each micro-colony, one for each time period, to determine the relationship between time and each of the three variables: syrup collection; pollen collection; weight gain. The slopes of the regressions for each time period were analysed for effects of pesticide exposure using a GLM.

## Results

### Worker mortality & weight change

Over the 5 week study period, a total of 6 worker bees died with at least one death per treatment group. No one micro-colony had more than 1 death and the total number of deaths was too few for further analysis. All workers lost weight during the study (fig. 1). However, workers in colonies that were exposed to pesticides lost significantly more weight than those that received uncontaminated food (GLM:  $\chi^2=5.10$ ,  $df=1$ ,  $p=0.02$ ). There was no significant effect of pollen diet on worker weight change (GLM:  $\chi^2=0.69$ ,  $df=1$ ,  $p=0.41$ ), nor was there any significant interaction between pesticide exposure and pollen diet (GLM:  $\chi^2=0.01$ ,  $df=1$ ,  $p=0.93$ ).

### Micro-colony growth, reproductive success & male quality

The amount that micro-colonies grew was significantly affected by both pollen diet and exposure to pesticides, with colonies receiving polyfloral pollen without pesticides performing best, and those receiving monofloral pollen contaminated with pesticides performing worst (Pollen, GLM:  $\chi^2=9.37$ ,  $df=1$ ,  $p=0.002$ ; pesticide, GLM:  $\chi^2=6.32$ ,  $df=1$ ,  $p=0.012$ ). There was no significant interaction between



the two factors (GLM:  $\chi^2=1.57$ ,  $df=1$ ,  $p=0.210$ ). Micro-colonies that received a monofloral diet grew on average 15.5% less than those that received a polyfloral pollen. Furthermore, micro-colonies that received uncontaminated syrup and pollen grew on average 15.6% more than micro-colonies that were exposed to pesticides (fig. 2 & fig. 3). Comparing the rate of weight gain during and after exposure, micro-colonies that received pesticides grew faster once they were no longer being exposed to pesticides than micro-colonies that received uncontaminated food throughout the experiment (GLM:  $\chi^2=6.44$ ,  $df=1$ ,  $p=0.011$ ) (fig. 2).

The number of males produced was significantly affected by the treatment applied (Kruskal-Wallis:  $\chi^2=21.27$ ,  $df=3$ ,  $p<0.001$ ) (fig. 4A). Micro-colonies that received monofloral pollen produced significantly fewer males than those fed polyfloral pollen (Mann-Whitney U:  $U=44$ ,  $df=38$ ,  $p<0.001$ ). Although pesticide treated micro-colonies produced fewer males than those that received uncontaminated food, this difference was not significant (Mann-Whitney U:  $U=142$ ,  $df=38$ ,  $p=0.108$ ). There was also a significant effect of pollen diet on the average number of brood per treatment group (GLM:  $\chi^2=18.78$ ,  $df=1$ ,  $p<0.001$ ). Micro-colonies that received monofloral pollen produced on average 32 fewer larvae and pupae and this represented a 40% reduction compared to polyfloral groups (fig. 4B).

Furthermore males from micro-colonies that were supplied with monofloral pollen were on average 0.05 g lighter than those fed polyfloral pollen (GLM:  $\chi^2=29.5$ ,  $df=1$ ,  $p<0.001$ ). The weight of males was not significantly affected by pesticide exposure (GLM:  $\chi^2=0.50$ ,  $df=1$ ,  $p=0.481$ ) (fig. 5A.). The male thorax width was also significantly lower in micro-colonies given monofloral pollen compared to polyfloral pollen (GLM:  $\chi^2=20.1$ ,  $df=1$ ,  $p<0.001$ ). There was also a significant effect of pesticide exposure on male thorax width, with thoraxes being narrower in micro-colonies that were exposed to pesticides (GLM:  $\chi^2=5.57$ ,  $df=1$ ,  $p=0.018$ ). There was a marginally non-significant interaction between pesticide exposure and pollen diet on male thorax width (GLM:  $\chi^2=3.74$ ,  $df=1$ ,  $p=0.053$ ) (fig. 5B). The fat content of males, measures as a proportion of body weight, was significantly higher in micro-colonies fed on a polyfloral diet ( $F_{1,24} = 24.2$ ,  $p<0.001$ ) but there was no significant effect of pesticide contamination ( $F_{1,24} = 1.31$ ,  $p=0.264$ ), and no significant interaction between the two (fig. 6).

## Food collection

Micro-colonies that received monofloral pollen collected significantly less syrup than micro-colonies that received polyfloral pollen (GLM:  $\chi^2=7.42$ ,  $df=1$ ,  $p=0.006$ ) (fig. 7A). However, when controlling for variation in the amount of weight gained by micro-colonies (weight gain was added as a covariate to the GLM), this difference was non-significant (GLM:  $\chi^2=2.07$ ,  $df=1$ ,  $p=0.150$ ). There was no significant effect of pesticide exposure on the amount of syrup collected by micro-colonies, both

before and after controlling for micro-colony weight gain (before, GLM:  $\chi^2=1.06$ ,  $df=1$ ,  $p=0.304$ ; after, GLM:  $\chi^2=0.12$ ,  $df=1$ ,  $p=0.726$ ).

There was no significant effect of pollen diet on the amount of pollen collected by micro-colonies, both before and after controlling for the amount of weight gained (before, GLM:  $\chi^2<0.001$ ,  $df=1$ ,  $p=0.985$ ; after, GLM:  $\chi^2=2.88$ ,  $df=1$ ,  $p=0.090$ ). There was also no significant effect of pesticide exposure on the amount of pollen collected by micro-colonies, both before and after controlling for weight gain (before, GLM:  $\chi^2=3.27$ ,  $df=1$ ,  $p=0.070$ ; after, GLM:  $\chi^2=0.39$ ,  $df=1$ ,  $p=0.534$ ).

Micro-colonies that received pesticide contaminated food collected both pollen and syrup in significantly greater quantity in the period after exposure than micro-colonies that received non-contaminated food throughout (syrup, GLM:  $\chi^2=7.897$ ,  $df=1$ ,  $p=0.005$ ; pollen, GLM:  $\chi^2=6.441$ ,  $df=1$ ,  $p=0.011$ ).

The average amount of thiamethoxam removed from the feeders per worker over the experimental period was 27.22 ng (comprising 2.26 ng from pollen and 24.97 ng from syrup) in the monofloral micro-colonies and 29.25 ng (comprising 2.07 ng from pollen and 27.18 ng from syrup) in the polyfloral micro-colonies.

## Discussion

Both managed and wild pollinators are increasingly exposed to a wide range of threats, with many different pressures implicated in driving losses in their stocks at a global scale (Brown & Paxton, 2009; Potts et al., 2010; Goulson et al., 2015). Intensification in agricultural practices results not only in large-scale habitat loss, fragmentation and degradation, but also homogenises landscapes, reducing the diversity of floral resources, and often exposing bees to cocktails of harmful agrochemicals (Potts et al., 2010; Goulson et al., 2015). Here, we investigate for the first time the combined effects of varying diet quality and exposure to a neonicotinoid pesticide, thiamethoxam, on bee colony performance.

Consistent with the findings of previous studies, neither a monotonous diet nor exposure to thiamethoxam at field-realistic levels (3.5 ppb) were sufficient to cause any significant worker mortality during the period studied. Reductions in worker survivorship relating to pollen diet have only been documented when workers are fed solely on syrup and deprived of pollen altogether (Duchateau & Velthuis, 1989; Génissel et al., 2002; Smeets & Duchateau, 2003). Furthermore, studies relating to the effects of thiamethoxam in bumblebees have only detected significant reductions in worker life expectancy at concentrations above 100 ppb (Mommaerts et al., 2010; Laycock et al., 2014), nearly 30 times higher than the dosage applied to pollen and syrup in this study.

Although there were no lethal effects, we detected a variety of sub-lethal effects in this study. For example, micro-colonies that received monofloral pollen gained less weight (figs. 2 & 3), and exhibited lower reproductive effort; monofloral micro-colonies produced fewer males (fig. 4A) and had fewer larvae and pupae than micro-colonies that received a polyfloral diet (fig. 4B). Not only was the total reproductive output of monofloral micro-colonies lower, the quality of drones produced was also significantly reduced, with males being lighter (fig. 5A) and smaller (fig. 5B) and with lower lipid content (fig. 6). Our findings are broadly in agreement with those of previous studies which generally find that colonies perform comparatively poorly when fed a monofloral diet (Génissel et al., 2002; Tasei & Aupinel, 2008; Baloglu & Gurel, 2015). However, caution is needed in interpreting our results. The protein content of the monofloral pollen was 24% lower than the polyfloral pollen blend; studies by Greenberg (1982) and Regali & Rasmont (1995) have shown that higher pollen protein consumption can increase the size of bees. Alternatively, differences between treatments may be a consequence of differences in other nutritive properties, such as the composition of amino acids or sterols. A recent study by Vanderplanck et al. (2014) investigating how pollen chemistry of five different monofloral pollens affected the development of bumblebee colonies found that the most important factors determining pollen performance were the polypeptides/total amino acids concentration and sterol composition; the two pollens that performed the best contained high concentrations of polypeptides/total amino acids and the sterol 24-methylenecholesterol.

Some studies have indicated that monofloral pollens of better quality can produce levels of colony performance comparable to polyfloral blends (Génissel et al., 2002; Baloglu & Gurel, 2015) and *Cistus* pollen specifically has been shown to perform badly compared to other monofloral pollens (Tasei & Aupinel, 2008a; Baloglu & Gurel, 2015; Moerman et al., 2015); Moerman et al. (2015) demonstrated that *Cistus* fed colonies grew slower than colonies fed either *Salix* or *Actinidia deliciosa* pollen and that this was down to *Cistus* pollen's lower amino acid concentration. Overall, it is clear that pollen diet has profound implications for bee colonies, but further work is required before we can draw general conclusions as to what constitutes a healthy diet for bees.

Pesticide exposure had fewer detectable effects than diet. Micro-colonies that received contaminated food gained less weight than micro-colonies that received uncontaminated food (figs. 2 & 3), but their total reproductive output was comparable to that of uncontaminated micro-colonies (fig. 4). Our findings seem to be largely in accordance with those of Elston et al. (2013), who detected reductions in micro-colony performance at thiamethoxam doses within a field-realistic range. Conversely, both Mommaerts et al. (2010) and Laycock et al. (2014) did not detect any effects, even when doses were 10 ppb which is nearer the likely upper limit for average field exposure (Botias et al. 2015). As previously mentioned however, both of these studies only dosed

343 dietary syrup and not pollen, and so undoubtedly underestimate extent of exposure that would be  
344 experienced by bees in the wild. Moreover, these studies only investigated quantitative measures of  
345 reproduction (i.e. numbers of brood) and not qualitative measures such as offspring size and lipid  
346 content. We found that pesticide exposure decreased the size of males (as measured by thorax  
347 width). As previous work has found that micro-colonies are representative analogues of whole  
348 colonies (Tasei & Aupinel 2008b), it is plausible that similar patterns would also be seen in the  
349 production of new workers and queens. As smaller queens and males generally experience lower  
350 reproductive success, with smaller queens being less likely to survive winter hibernation and smaller  
351 males less likely to mate (Beekman et al., 1998a; Beekman 1998b; Amin et al., 2012; Vanderplanck et  
352 al., 2014), this could have important consequences for reproductive success and fitness at the  
353 population level.

354         It should also be noted that, as food was provided directly within micro-colony boxes, there  
355 was no need for bees to forage. One of the main influences of neonicotinoids on bees is through  
356 their impairment of foraging behaviour and homing ability (Yang et al., 2008; Schneider et al., 2012;  
357 Feltham et al. 2014), and Mommaerts et al (2010) demonstrated that bumblebees are up to 10  
358 times more sensitive to imidacloprid when they have to forage compared to when food is provided  
359 directly. Additionally, wild bees may be exposed to a wide range of additional stressors, such as  
360 other chemicals including EBI fungicides and infection from pathogens, both of which have been  
361 shown to amplify the adverse effects of neonicotinoids (Pilling & Jepson, 1993; Schmuck et al., 2003;  
362 Di Prisco et al., 2013; Sgolastra et al., 2016). Thus we might expect greater effects of pesticides on  
363 bee colonies under more natural settings.

364         When micro-colony weight gain was added as a covariate, there was no significant  
365 difference in how much syrup was collected by micro-colonies, suggesting that the size of the micro-  
366 colonies was the most significant factor in explaining the variation in syrup collection (fig. 7A.). In  
367 contrast to syrup collection, pollen quality did not significantly affect pollen collection. This is in  
368 accordance with Mommaerts et al. (2010) and Vanderplanck et al. (2014) suggesting that workers do  
369 not adjust their consumption or larval provisions when pollen is nutritionally poor. There is evidence  
370 that bumblebees can assess the chemical quality of pollen, enabling them to select pollen of  
371 superior quality (Robertson et al., 1999; Hanley et al., 2008; Kitaoka & Nieh, 2009; Leonhardt &  
372 Blüthgen, 2012), yet it does not seem that they compensate when collecting low quality pollen by  
373 collecting more.

374         We detected that in both groups that were exposed to pesticides, micro-colonies collected  
375 pollen and nectar more quickly when they had been provided with uncontaminated food, suggesting  
376 that there may have been an anti-feedant effect imposed by thiamethoxam. Whilst it has been  
377 shown that anti-feedant properties of neonicotinoids can lead to reduced reproduction (Gill et al.,

2012; Laycock et al., 2012; Elston et al., 2013; Laycock et al., 2014), these anti-feedant effects have only been noticed when doses were in excess of 10 ng/g. As the reproductive effort of colonies that were exposed to pesticides was comparable to those received uncontaminated food (fig. 4), it seems that micro-colonies were able to compensate for anti-feedant effects by eating more and growing more quickly once no longer exposed to thiamethoxam (fig. 2).

Overall, our findings suggest that both dietary pollen quality and exposure to the neonicotinoid thiamethoxam have multiple, measurable adverse effects on bumblebee micro-colonies, though there were no strong interactions between the two stressors. Although micro-colonies are regarded as being good proxies for whole colonies, it would be informative to investigate the effects of these factors using whole colonies in more realistic field settings where other stressors are present and where bees have to forage to collect food.

## References

- Alaux, C., Brunet, J. L., Dussaubat, C., Mondet, F., Tchamitchan, S., Cousin, M., ... & Le Conte, Y. (2010). Interactions between *Nosema* microspores and a neonicotinoid weaken honeybees (*Apis mellifera*). *Environmental Microbiology*, 12(3), 774-782.
- Amin, M. R., Bussière, L. F., & Goulson, D. (2012). Effects of male age and size on mating success in the bumblebee *Bombus terrestris*. *Journal of Insect Behavior*, 25(4), 362-374.
- Arnold, G., Pistorius, J., Steeger, T., & Thompson, H. (2012). Statement on the findings in recent studies investigating sub-lethal effects in bees of some neonicotinoids in consideration of the uses currently authorised in Europe. *EFSA J*, 10, 2752-2779.
- Baloglu, G. H., & Gurel, F. (2015). The effects of pollen protein content on colony development of the bumblebee, *Bombus terrestris* L. *Journal of Apicultural Science*, 59(1), 83-88.
- Beekman, M., Van Stratum, P., & Lingeman, R. (1998a). Diapause survival and post-diapause performance in bumblebee queens (*Bombus terrestris*). *Entomologia Experimentalis et Applicata*, 89(3), 207-214.
- Beekman, M., & Van Stratum, P. (1998b). Bumblebee sex ratios: why do bumblebees produce so many males? *Proceedings of the Royal Society of London B: Biological Sciences*, 265(1405), 1535-1543.
- Biesmeijer, J.C., Roberts, S.P.M., Reemer, M., Ohlemüller, R., Edwards, M., Peeters, T., Schaffers, A.P., Potts, S.G., Kleukers, R., Thomas, C.D. and Settele, J., (2006). Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science*, 313(5785), 351-354.
- Blacquièrè, T., Smagghe, G., Van Gestel, C. A., & Mommaerts, V. (2012). Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment. *Ecotoxicology*, 21(4), 973-992.

412 Botías, C., David, A., Horwood, J., Abdul-Sada, A., Nicholls, E., Hill, E. and Goulson, D., (2015).  
 413 Neonicotinoid residues in wildflowers, a potential route of chronic exposure for  
 414 bees. *Environmental Science & Technology*, 49(21), 12731-12740.

415 Bradford, M.M., (1976). A rapid and sensitive method for the quantitation of microgram quantities of  
 416 protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1-2), 248-  
 417 254.

418 Brown, M. J. F., Loosli, R., & Schmid-Hempel, P. (2000). Condition-dependent expression of virulence  
 419 in a trypanosome infecting bumblebees. *Oikos*, 91(3), 421-427.

420 Brown, M. J., & Paxton, R. J. (2009). The conservation of bees: a global perspective. *Apidologie*, 40(3),  
 421 410-416.

422 Cameron, S.A., Lozier, J.D., Strange, J.P., Koch, J.B., Cordes, N., Solter, L.F. and Griswold, T.L., (2011).  
 423 Patterns of widespread decline in North American bumble bees. *Proceedings of the National*  
 424 *Academy of Sciences*, 108(2), 662-667.

425 Carvell, C. (2002). Habitat use and conservation of bumblebees (*Bombus spp.*) under different  
 426 grassland management regimes. *Biological Conservation*, 103(1), 33-49.

427 Carvell, C., Roy, D. B., Smart, S. M., Pywell, R. F., Preston, C. D., & Goulson, D. (2006). Declines in  
 428 forage availability for bumblebees at a national scale. *Biological Conservation*, 132(4), 481-  
 429 489.

430 Di Pasquale, G., Salignon, M., Le Conte, Y., Belzunces, L. P., Decourtye, A., Kretzschmar, A., ... & Alaux,  
 431 C. (2013). Influence of pollen nutrition on honey bee health: do pollen quality and diversity  
 432 matter? *PloS one*, 8(8), e72016.

433 Di Prisco, G., Cavaliere, V., Annoscia, D., Varricchio, P., Caprio, E., Nazzi, F., ... & Pennacchio, F. (2013).  
 434 Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a  
 435 viral pathogen in honey bees. *Proceedings of the National Academy of Sciences*, 110(46),  
 436 18466-18471.

437 Dively, G. P., & Kamel, A. (2012). Insecticide residues in pollen and nectar of a cucurbit crop and their  
 438 potential exposure to pollinators. *Journal of Agricultural and Food Chemistry*, 60(18), 4449-  
 439 4456.

440 Duchateau, M. J., & Velthuis, H. H. W. (1989). Ovarian development and egg laying in workers of  
 441 *Bombus terrestris*. *Entomologia Experimentalis et Applicata*, 51(3), 199-213.

442 EFSA. (2013). EFSA Guidance Document on the risk assessment of plant protection products on bees  
 443 (*Apis mellifera*, *Bombus spp.* and solitary bees).

444 Elbert, A., Haas, M., Springer, B., Thielert, W., & Nauen, R. (2008). Applied aspects of neonicotinoid  
 445 uses in crop protection. *Pest Management Science*, 64(11), 1099-1105.

446 Elston, C., Thompson, H. M., & Walters, K. F. (2013). Sub-lethal effects of thiamethoxam, a  
 447 neonicotinoid pesticide, and propiconazole, a DMI fungicide, on colony initiation in  
 448 bumblebee (*Bombus terrestris*) micro-colonies. *Apidologie*, 44(5), 563-574.

449 Feltham, H., Park, K.J. and Goulson, D. 2014. Field Realistic Doses of Pesticide Imidacloprid Reduce  
 450 Bumblebee Pollen Foraging Efficiency. *Ecotoxicology* 23: 317-323.

451 Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., ... & Helkowski, J. H.  
 452 (2005). Global consequences of land use. *Science*, 309(5734), 570-574.

453 Gallai, N., Salles, J.M., Settele, J. and Vaissière, B.E., (2009). Economic valuation of the vulnerability of  
 454 world agriculture confronted with pollinator decline. *Ecological Economics*, 68(3), 810-821.

455 Génissel, A., Aupinel, P., Bressac, C., Tasei, J. N., & Chevrier, C. (2002). Influence of pollen origin on  
 456 performance of *Bombus terrestris* micro-colonies. *Entomologia Experimentalis et*  
 457 *Applicata*, 104(2-3), 329-336.

458 Gill, R. J., Ramos-Rodriguez, O., & Raine, N. E. (2012). Combined pesticide exposure severely affects  
 459 individual-and colony-level traits in bees. *Nature*, 491(7422), 105-108.

460 Godfray, H. C. J., Blacquiere, T., Field, L. M., Hails, R. S., Petrokofsky, G., Potts, S. G., ... & McLean, A. R.  
 461 (2014). A restatement of the natural science evidence base concerning neonicotinoid  
 462 insecticides and insect pollinators. *Proceedings of the Royal Society of London B: Biological*  
 463 *Sciences*, 281(1786), 20140558.

464 Goulson, D., Lye, G.C. and Darvill, B., (2008). Decline and conservation of bumble bees. *Annu. Rev.*  
 465 *Entomol.*, 53, 191-208.

466 Goulson, D. (2013). Review: An overview of the environmental risks posed by neonicotinoid  
 467 insecticides. *Journal of Applied Ecology*, 50(4), 977-987.

468 Goulson D. (2015). Neonicotinoids impact bumblebee colony fitness in the field; a reanalysis of the  
 469 UK's Food & Environment Research Agency 2012 experiment. *PeerJ* 3:e854.

470 Goulson, D., Nicholls, E., Botías, C. and Rotheray, E.L., (2015). Bee declines driven by combined stress  
 471 from parasites, pesticides, and lack of flowers. *Science*, 347(6229), p.1255957

472 Graystock, P., Jones, J.C., Pamminger, T., Parkinson, J.F., Norman, V., Blane, E.J., Rothstein, L. et al.  
 473 2016. Hygienic food to reduce pathogen risk to bumblebees. *Journal of Invertebrate Pathology*  
 474 136: 68-73.

475 Greenberg, L. (1982). Year-round culturing and productivity of a sweat bee, *Lasioglossum zephyrum*  
 476 (Hymenoptera: Halictidae). *Journal of the Kansas Entomological Society*, 13-22.

477 Grixti, J.C., Wong, L.T., Cameron, S.A. and Favret, C., (2009). Decline of bumble bees (*Bombus*) in the  
 478 North American Midwest. *Biological Conservation*, 142(1), 75-84.

479 Hanley, M. E., Franco, M., Pichon, S., Darvill, B., & Goulson, D. (2008). Breeding system, pollinator  
480 choice and variation in pollen quality in British herbaceous plants. *Functional Ecology*, 22(4),  
481 592-598.

482 Henry, M., Beguin, M., Requier, F., Rollin, O., Odoux, J. F., Aupinel, P., ... & Decourtye, A. (2012). A  
483 common pesticide decreases foraging success and survival in honey bees. *Science*, 336(6079),  
484 348-350.

485 Higes, M., Martín-Hernández, R., Garrido-Bailón, E., García-Palencia, P. and Meana, A., (2008).  
486 Detection of infective *Nosema ceranae* (Microsporidia) spores in corbicular pollen of forager  
487 honeybees. *Journal of Invertebrate Pathology*, 97(1), 76-78.

488 Howard, D. C., Watkins, J. W., Clarke, R. T., Barnett, C. L., & Stark, G. J. (2003). Estimating the extent  
489 and change in broad habitats in Great Britain. *Journal of Environmental Management*, 67(3),  
490 219-227.

491 Human, H., Nicolson, S. W., Strauss, K., Pirk, C. W. W., & Dietemann, V. (2007). Influence of pollen  
492 quality on ovarian development in honeybee workers (*Apis mellifera scutellata*). *Journal of*  
493 *Insect Physiology*, 53(7), 649-655.

494 James, R. R., & Xu, J. (2012). Mechanisms by which pesticides affect insect immunity. *Journal of*  
495 *Invertebrate Pathology*, 109(2), 175-182.

496 Junjie, F., Weiqiao, S. & Jianhua, Z. (1998). Effect of radiation sterilization on pine pollen storage  
497 quality. *Proceedings of the 7<sup>th</sup> International Working Conference on Stored-product Protection*,  
498 2, Sichuan Publishing House of Science and Technology, Chengdu, China, pp. 1701-1704. (ISBN  
499 7536440987)

500 Kitaoka, T. K., & Nieh, J. C. (2009). Bumble bee pollen foraging regulation: role of pollen quality,  
501 storage levels, and odor. *Behav Ecol Sociobiol*, 63, 501-510

502 Kleijn, D., & Raemakers, I. (2008). A retrospective analysis of pollen host plant use by stable and  
503 declining bumble bee species. *Ecology*, 89(7), 1811-1823.

504 Kollmeyer, W. D., Flattum, R. F., Foster, J. P., Powell, J. E., Schroeder, M. E., & Soloway, S. B. (1999).  
505 Discovery of the nitromethylene heterocycle insecticides. In: *Nicotinoid insecticides and the*  
506 *nicotinic acetylcholine receptor*, Yamamoto et al. (ed.), Springer-Verlag, Tokyo, 71-89.

507 Kosior, A., Celary, W., Olejniczak, P., Fijal, J., Krol, W., Solarz, W. and Plonka, P., (2007). The decline of  
508 the bumble bees and cuckoo bees (Hymenoptera: Apidae: Bombini) of Western and Central  
509 Europe. *Oryx*, 41(01), 79-88.

510 Kremen, C., Williams, N. M., Aizen, M. A., Gemmill-Herren, B., LeBuhn, G., Minckley, R., ... & Winfree,  
511 R. (2007). Pollination and other ecosystem services produced by mobile organisms: a  
512 conceptual framework for the effects of land-use change. *Ecology Letters*, 10(4), 299-314.



513 Laycock, I., Lenthall, K. M., Barratt, A. T., & Cresswell, J. E. (2012). Effects of imidacloprid, a  
 514 neonicotinoid pesticide, on reproduction in worker bumble bees (*Bombus*  
 515 *terrestris*). *Ecotoxicology*, 21(7), 1937-1945.

516 Laycock, I., Cotterell, K. C., O'Shea-Wheller, T. A., & Cresswell, J. E. (2014). Effects of the neonicotinoid  
 517 pesticide thiamethoxam at field-realistic levels on micro-colonies of *Bombus terrestris* worker  
 518 bumble bees. *Ecotoxicology and Environmental Safety*, 100, 153-158.

519 Moerman, R., Vanderplanck, M., Roger, N., Declèves, S., Wathelet, B., Rasmont, P., ... & Michez, D.  
 520 (2015). Growth rate of bumblebee larvae is related to pollen amino acids. *Journal of Economic*  
 521 *Entomology*, tov279.

522 Mommaerts, V., Reynders, S., Boulet, J., Besard, L., Sterk, G., & Smagghe, G. (2010). Risk assessment  
 523 for side-effects of neonicotinoids against bumblebees with and without impairing foraging  
 524 behavior. *Ecotoxicology*, 19(1), 207-215.

525 Osborne, J. L., Martin, A. P., Carreck, N. L., Swain, J. L., Knight, M. E., Goulson, D., ... & Sanderson, R. A.  
 526 (2008). Bumblebee flight distances in relation to the forage landscape. *Journal of Animal*  
 527 *Ecology*, 77(2), 406-415.

528 Pettis, J. S., Johnson, J., & Dively, G. (2012). Pesticide exposure in honey bees results in increased  
 529 levels of the gut pathogen *Nosema*. *Naturwissenschaften*, 99(2), 153-158.

530 Pilling, E. D., & Jepson, P. C. (1993). Synergism between EBI fungicides and a pyrethroid insecticide in  
 531 the honeybee (*Apis mellifera*). *Pesticide Science*, 39(4), 293-297.

532 Potts, S.G., Roberts, S.P., Dean, R., Marris, G., Brown, M.A., Jones, R., Neumann, P. and Settele, J.,  
 533 (2010). Declines of managed honey bees and beekeepers in Europe. *Journal of Apicultural*  
 534 *Research*, 49(1), 15-22.

535 Rasmont, P., Pauly, A., Terzo, M., Patiny, S., Michez, D., Iserbyt, S., ... & Haubruge, E. (2005). The  
 536 survey of wild bees (Hymenoptera, Apoidea) in Belgium and France. *Food and Agriculture*  
 537 *Organisation, Rome*, 18.

538 Regali, A., & Rasmont, P. (1995). Nouvelles méthodes de test pour l'évaluation du régime alimentaire  
 539 chez des colonies orphelines de *Bombus terrestris* (L) (Hymenoptera,  
 540 Apidae). *Apidologie*, 26(4), 273-281.

541 Renzi, M.T., Rodríguez-Gasol, N., Medrzycki, P., Porrini, C., Martini, A., Burgio, G., Maini, S., Sgolastra,  
 542 F. 2016. Combined effect of pollen quality and thiamethoxam on hypopharyngeal gland  
 543 development and protein content in *Apis mellifera*. *Apidologie*, in press.

544 Ricketts, T. H., Regetz, J., Steffan - Dewenter, I., Cunningham, S. A., Kremen, C., Bogdanski, A., ... &  
 545 Morandin, L. A. (2008). Landscape effects on crop pollination services: are there general  
 546 patterns? *Ecology letters*, 11(5), 499-515.

547 Roulston, T. H., Cane, J. H., & Buchmann, S. L. (2000). What governs protein content of pollen:  
 548 pollinator preferences, pollen–pistil interactions, or phylogeny? *Ecological Monographs*, 70(4),  
 549 617–643.

550 Schmuck, R., Stadler, T., & Schmidt, H. W. (2003). Field relevance of a synergistic effect observed in  
 551 the laboratory between an EBI fungicide and a chloronicotinyl insecticide in the honeybee  
 552 (*Apis mellifera* L, Hymenoptera). *Pest Management Science*, 59(3), 279–286.

553 Schneider, C. W., Tautz, J., Grünewald, B., & Fuchs, S. (2012). RFID tracking of sublethal effects of two  
 554 neonicotinoid insecticides on the foraging behavior of *Apis mellifera*. *PloS one*, 7(1), e30023.

555 Sgolastra, F., Medrzycki, P., Bortolotti, L., Renzi, M.T., Tosi, S., Bogo, G., Teper, D. et al. 2016.  
 556 Synergistic interactions between pesticides in three bee species. *Pest Management Science*, in  
 557 press.

558 Singh, R., Levitt, A.L., Rajotte, E.G., Holmes, E.C., Ostiguy, N., Lipkin, W.I., Toth, A.L. and Cox-Foster,  
 559 D.L., (2010). RNA viruses in hymenopteran pollinators: evidence of inter-taxa virus  
 560 transmission via pollen and potential impact on non-*Apis* hymenopteran species. *PLoS*  
 561 *One*, 5(12), p.e14357.

562 Smeets, P., & Duchateau, M. (2003). Longevity of *Bombus terrestris* workers (Hymenoptera: Apidae) in  
 563 relation to pollen availability, in the absence of foraging. *Apidologie*, 34(4), 333–337.

564 Stoner, K. A., & Eitzer, B. D. (2012). Movement of soil-applied imidacloprid and thiamethoxam into  
 565 nectar and pollen of squash (*Cucurbita pepo*). *PLoS One*, 7(6), e39114.

566 Tasei, J. N., Lerin, J., & Ripault, G. (2000). Sub-lethal effects of imidacloprid on bumblebees, *Bombus*  
 567 *terrestris* (Hymenoptera: Apidae), during a laboratory feeding test. *Pest Management*  
 568 *Science*, 56(9), 784–788.

569 Tasei, J. N., & Aupinel, P. (2008a). Nutritive value of 15 single pollens and pollen mixes tested on  
 570 larvae produced by bumblebee workers (*Bombus terrestris*, Hymenoptera:  
 571 Apidae). *Apidologie*, 39(4), 397–409.

572 Tasei, J. N., & Aupinel, P. (2008b). Validation of a method using queenless *Bombus terrestris* micro-  
 573 colonies for testing the nutritive value of commercial pollen mixes by comparison with  
 574 queenright colonies. *Journal of Economic Entomology*, 101(6), 1737–1742.

575 Thermo Scientific, (2010). *PROTOCOL NanoDrop 2000/2000c Bradford Protein Assay*. [Online].  
 576 Available at: [http://www.nanodrop.com/Library/T084-NanoDrop-2000-2000c-Bradford-](http://www.nanodrop.com/Library/T084-NanoDrop-2000-2000c-Bradford-Protocol.pdf)  
 577 [Protocol.pdf](http://www.nanodrop.com/Library/T084-NanoDrop-2000-2000c-Bradford-Protocol.pdf) [accessed: 23/02/2016]

578 Vanderplanck, M., Moerman, R., Rasmont, P., Lognay, G., Wathelet, B., Wattiez, R., & Michez, D.  
 579 (2014). How does pollen chemistry impact development and feeding behaviour of polylectic  
 580 bees? *PloS one*, 9(1), e86209.

581 Westphal, C., Steffan-Dewenter, I., & Tscharntke, T. (2003). Mass flowering crops enhance pollinator  
582 densities at a landscape scale. *Ecology Letters*, 6(11), 961-965.

583 Whitehorn, P. R., O'Connor, S., Wackers, F. L., & Goulson, D. (2012). Neonicotinoid pesticide reduces  
584 bumble bee colony growth and queen production. *Science*, 336(6079), 351-352.

585 Williams, P.H. and Osborne, J.L., (2009). Bumblebee vulnerability and conservation world-  
586 wide. *Apidologie*, 40(3), 367-387.


587 Winfree, R., Aguilar, R., Vázquez, D. P., LeBuhn, G., & Aizen, M. A. (2009). A meta-analysis of bees'  
588 responses to anthropogenic disturbance. *Ecology*, 90(8), 2068-2076.

589 Winfree, R., (2010). The conservation and restoration of wild bees. *Annals of the New York Academy of*  
590 *Sciences*, 1195(1), 169-197.

591 Xie, Z., Williams, P.H. and Tang, Y., (2008). The effect of grazing on bumblebees in the high rangelands  
592 of the eastern Tibetan Plateau of Sichuan. *Journal of Insect Conservation*, 12(6), 695-703.

593 Yang, E. C., Chuang, Y. C., Chen, Y. L., & Chang, L. H. (2008). Abnormal foraging behavior induced by  
594 sublethal dosage of imidacloprid in the honey bee (Hymenoptera: Apidae). *Journal of*  
595 *economic entomology*, 101(6), 1743-1748.


596



**Figure 1.** The average worker weight change per individual bee for each treatment group across the 5 week study period (median and interquartile range). All workers lost weight. M = monofloral, P = polyfloral, U = uncontaminated, C = contaminated with pesticide.


597

598




**Figure 2.** The average cumulative weight gain per micro-colony of treatment groups through time. The error bars show the standard error of all the microcolonies in each group. The vertical dashed line indicates the periods during and after pesticide exposure. Uncontaminated pollen and syrup was provided to all groups from day 17 onwards.

600




**Figure 3. Total weight gain.** The average weight gain of microcolonies from each treatment group at the end of the 5 weeks. M = monofloral, P = polyfloral, U = uncontaminated, C = contaminated with pesticide.

601




**Figure 4. A.** The average number of males produced by microcolonies from each group throughout the 5 week study. **B.** The average number of brood per micro-colony from each treatment group. M = monofloral, P = polyfloral, U = uncontaminated, C = contaminated with pesticide.




**Figure 5. A.** The average weight of individual males produced by each treatment group. **B.** The average thorax width of males per group. M = monofloral, P = polyfloral, U = uncontaminated, C = contaminated with pesticide.





**Figure 6. A.** The average fat content of male offspring, expressed as a proportion of body weight, from each treatment group. M = monofloral, P = polyfloral, U = uncontaminated, C = contaminated with pesticide.



**Figure 7. A.** The average amount of syrup collected by microcolonies of each treatment group across the 5 week study period. **B.** The average amount of pollen collected by microcolonies of different treatment groups across the 5 weeks.